

The Effect of Methanol Leaf Extract of *Rauwolfia vomitoria* on Hepatic Markers of Chloroform Intoxicated Albino Wistar Rats

Ugwu Okechukwu Paul Chima, Agu Chibuike Cyprian and Okon Michael Ben
Department of Biochemistry, Faculty of Science, Ebonyi State University,
Abakaliki.

ABSTRACT

Medicinal plants are the richest sources for drugs of traditional systems of medicine, modern medicines, nutraceuticals, food supplements, folk medicines, pharmaceutical intermediates and chemical entities for synthetic drugs. They are one of the most important sources of new chemical compounds with potential therapeutic effects. *Rauwolfia vomitoria* is one of the medicinal plants that have served all through the ages as the mainstay in the treatment and preservation of human health. This research was designed to determine the effects of methanol leaf extract of *Rauwolfia vomitoria* on hepatic markers of (ALT, ALP, AST, GGT and bilirubin) on chloroform intoxicated wistar albino rats. All chemicals and reagents used in this research were of analytical grade. The results indicated that ALT, ALP, AST, GGT activities and bilirubin level decreased significantly ($p < 0.05$) in groups 2, 3 and 4 rats treated with graded doses of 100 mg/kg, 200mg/kg and 300mg/kg b.w of methanol leaf extract of *Rauwolfia vomitoria* when compared with the positive control rats treated with 0.5 ml of normal saline. Treatment with graded doses of methanol leaf extract produced significant decrease ($p < 0.05$) in ALT, ALP, AST, GGT activities and bilirubin level when compared with the positive control group. In conclusion, the results from this research indicated that methanol leaf extract of *Rauwolfia vomitoria* can decrease liver markers in chloroform intoxicated rats as shown in the treated groups compared with the untreated control (Positive control). As a result methanol leaf extract of *Rauwolfia vomitoria* can be used to ameliorate hepatic associated diseases.

Keywords: *Rauwolfia Vomitoria*, Hepatic Markers, Chloroform.

INTRODUCTION

Medicinal plants are the richest bioresource of drugs of traditional systems of medicine, modern medicines, nutraceuticals, food supplements, folk medicines, pharmaceutical intermediates, and chemical entities for synthetic drugs [1,2,3,4,5]. They are one of the most important sources of new chemical compounds with potential therapeutic effects [6,7]. [8,,9] has advocated traditional medicine as safe remedies for ailments of both microbial and non-microbial origin. Plants are major sources of therapeutic compounds and are the essential foundation of medicine since prehistoric time [10,11,12]. Plants

synthesize thousands of chemical compounds possessing different properties like defense against insects, bacteria, fungi, diseases and herbivorous mammals [13,14,15]. Herbal and natural products have been used in folk medicine for centuries throughout the world. Some Indian medicines like Ayurveda, Sindha and Unani entirely and homeopathy to some extent, depend on plant materials or their derivatives for treating human diseases [16,17]. Medicinal plants are widely used in non-industrialized societies, mainly because they are readily available and cheaper than modern medicines [18,19]. Medicinal plants

have been discovered and used in traditional medicine practices since prehistoric times [20]. There has been renewed interest in screening higher plants for novel biologically active compounds, particularly those that effectively intervene in human ailments in the field of chronic diseases [21]. Currently, research is focused on the isolation of pharmacologically active compounds from natural sources in the area of those diseases where presently available drugs are not so effective [22,23]. Also herbal medicines are experiencing greater resurgence as many people are turning their attention from modern drugs toward parallel herbal systems which are also known as alternative medicine. Plants have been used for centuries as a remedy for human diseases because they possess phytochemicals of therapeutic values [24,25,26]. The Indian Traditional medicine like Ayurveda, Siddha and Unani are predominantly based on the use of plant materials [27,28,29]. Herbal drugs have gained importance and popularity in recent years because of their safety, efficacy and cost effectiveness. The association of medical plants with other plants in their habitat also influences their medicinal values in some cases [30]. One of the important and well documented uses of plant-products is their use as antioxidant [31,32]. Hence there is an ever increasing need for health safety in the society filled with toxicants [32]. In spite of

tremendous strides in modern medicine, in 2004, the U.S. National Centre for complementary and Alternative Medicine of the National Institutes of Health began funding clinical trials into the effectiveness of herbal medicine [9]. For this reason, various medicinal plants have been studied using modern scientific approaches which have shown that due to various biological components, many of these medicinal plants possess a number of properties such as anti-diabetic, antioxidant, anticancer and anti-inflammatory effects, etc. and can be used to treat a wide range of various diseases [14]. The medicinal properties of plants are due to the presence of certain specific substances, referred to as bioactive principles which may be stored in organs like roots, leaves, stem bark, fruits and seeds [16]. Many herbs have shown positive results in-vitro, animal models, or small-scale clinical tests [18]. Plants and their extracts have immense potentials for the management and treatment of wounds. The phytomedicine for wound healing are not only cheap and affordable, but are also safe as hypersensitive reactions are rarely encountered. These natural agents induce healing and regeneration of tissues by multiple mechanisms. However, according to [20], there is need for scientific validation, standardization and safety evaluation of plants of traditional medicine before recommendation for any therapeutic use.

Aim

The aim of this research was to determine the effects of methanol leaf extract of *Rauwolfia vomitoria* on hepatic markers

of chloroform intoxicated wistar albino rats.



Figure 1: The leaves of *Rauwolfia vomitoria*

MATERIALS AND METHODS

Methods

Preparation of Plant Material

The leaves of *Rauwolfia vomitoria* were collected, dried and milled to powder

using the grinding machine.

Extraction of Plant Material

A known quantity, 500g of ground leaves of *Rauwolfia vomitoria* were macerated in 1500ml of Methanol with thorough shaking at regular interval for 72h at room temperature (26-28°C). The resulting solution was filtered using Whatman No. 1

filter paper. The filtrates were concentrated using rotary evaporator to obtain slurry of the extract. The semi-pastry extract was stored in the refrigerator and used for the study.

Determination of Hepatic Markers

Hepatic Markers were done using the

standard methods of [20].

Experimental Design

Forty eight (48) Wistar albino rats were used in this study. They were randomly distributed into six (6) groups of 8 rats each. Oxidative stress was induced in the rats and this was performed by intraperitoneal injection of chloroform (100 mg/kg b/w). The rats were fed graded doses of ethanol extract of *Rauwolfia vomitoria* through oral intubation method. The groups and doses administered are summarized below:

Group 1: (Negative control rats without Chloroform intoxication): rats were treated with [0.5 ml of normal saline).

Group 2: (Chloroform intoxicated rats): rats were treated with (100 mg/kg b.w. of Methanol extract of *Rauwolfia vomitoria*).

Group 3: (Chloroform intoxicated rats): rats were treated with (200 mg/kg b.w. of Methanol extract of *Rauwolfia vomitoria*).

Group 4: (Chloroform intoxicated rats): rats were treated with (300 mg/kg b.w. of

Methanol extract of *Rauwolfia vomitoria*.

Group 5: (Positive control rats with Chloroform intoxication) were treated with (0.5 ml of normal saline).

Group 6: (Standard control rats with Chloroform intoxication) were treated with (5 mg/kg bodyweight of standard drug Chemiron)

Statistical Analysis

Results were expressed as mean± standard deviations where applicable. The data were subjected to one-way analysis of variance (ANOVA), followed by Post hoc

Duncan multiple comparison test using SPSS software version 21 and p < 0.05 was regarded as significant.

RESULTS

The results indicated that ALT, ALP, AST, GGT activities and bilirubin level decreased significantly (p<0.05) in groups 2, 3 and 4 rats treated with graded doses of 100 mg/kg, 200mg/kg and 300mg/kg b.w of methanol leaf extract of *Rauwolfia vomitoria* compared with the positive control rats treated with 0.5 ml of normal

saline as shown in figures 2, 3, 4, 5 and 6 Treatment with graded doses of methanol leaf extract produced significant decrease (p<0.05) in ALT, ALP, AST, GGT activities and bilirubin level compared with the positive control groups (figures 2, 3, 4, 5 and 6).




Fig 2: The ALT activity of rats treated with methanol extract of *Rauwolfia vomitoria*. Data are shown as mean ± standard deviation (n=4).




Fig 3: The ALP activity of rats treated with methanol extract of *Rauwolfia vomitoria*. Data are shown as mean \pm standard deviation (n=4).




Fig 4: The AST activity of rats treated with methanol extract of *Rauwolfia vomitoria*. Data are shown as mean \pm standard deviation (n=4).




Fig 5: The GGT activity of rats treated with methanol extract of *Rauwolfia vomitoria*. Data are shown as mean \pm standard deviation (n=4).




Fig 6: The Bilirubin Level of rats treated with methanol extract of *Rauwolfia vomitoria*. Data are shown as mean \pm standard deviation (n=4).

DISCUSSION

This study was designed to evaluate the ameliorative effect on the liver of chloroform intoxicated Wistar albino rats treated with graded doses of methanol leaf extract of *Rauwolfia vomitoria*. The results indicated that ALT, ALP, AST, GGT activities and bilirubin level decreased significantly ($p < 0.05$) in groups 2, 3 and 4 rats treated with graded doses of 100 mg/kg, 200mg/kg and 300mg/kg b.w of methanol leaf extract of *Rauwolfia vomitoria* compared with the positive control rats treated with 0.5 ml of normal saline as shown in figures 2, 3, 4, 5 and 6. Treatment with graded doses of methanol leaf extract produced significant decrease ($p < 0.05$) in ALT, ALP, AST, GGT activities and bilirubin level compared with the positive control groups (figures 2, 3, 4, 5

and 6). The decreased level in liver markers compared to the known standard drug in group 6 (Standard control) shows that the extract can be used to ameliorate hepatic damages in liver of chloroform intoxicated rats. This agrees with the work of [32] who obtained similar results on serum amino transferase and alkaline phosphatase activities of rats treated with *Rauwolfia vomitoria* Afzel (Apocynaceae) extract. The liver is the key site of metabolism of xenobiotics and also plays a role in synthesis of drugs. Liver malfunction will impair the function and metabolism of drugs and xenobiotics in the blood [32]. Aminotransferases are cytosolic enzymes widely distributed in tissues with highest concentration in liver and heart but with

ALT more specific to the liver and AST to the heart. Damage to the membrane architecture of cells due to exposure to toxicants will lead to their spillage into

CONCLUSION

In conclusion, the results from this research indicated that methanol leaf extract of *Rauwolfia vomitoria* can decrease liver markers in chloroform intoxicated rats as shown in the treated

www.iaajournals.org
blood. The increase in serum ALT therefore implies the toxicity of the extract on the liver [5].

groups compared with the untreated control (Positive control). As a result methanol leaf extract of *Rauwolfia vomitoria* can be used to ameliorate hepatic associated diseases.

REFERENCES

1. Aderemi, F. A. (2004). Effects of replacement of wheat bran with cassava root sieviate supplemented or unsupplemented with enzyme on the haematology and serum biochemistry of pullet chicks. *Tropical Journal of Animal Science*, 7, 147-153.
2. Afolabi, K. D., Akinsoyinu, A. O., Olajide, R., and Akinleye, S. B. (2010). Haematological parameters of the Nigerian local grower chickens fed varying dietary levels of palm kernel cake (p.247). Proceedings of 35th Annual Conference of Nigerian Society for Animal Production.
3. Agarwal, S. S. (2001). Development of hepatoprotective formulations from plant sources. In: Pharmacology and Therapeutics in the Millennium. Edited by Gupta SK, Narosa Publishing House, New Delhi, pp: 357-358.
4. Agency for Toxic Substances and Disease Registry (ATSDR) (2013). Toxicological Profile for Chloroform, US Department of Health and Human Services: Atlanta, US.
5. Akpanabiatu, M. I., Otitoju, O., Edet, E. E., Ndem, J. I., Uwah, A. F. and Ufot, U. F. (2012). Vitamin E Supplementation with *Rauwolfia Vomitoria* Root Bark Extract Improves Hematological Indices. *North American Journal of Medical Sciences*, 4(2): 86-89.
6. American Gastroenterological Association. (2002). American Gastroenterological Association Medical Position Statement: Evaluation of Liver Chemistry Tests. *Gastroenterology*.123(4):136 4-6 2.
7. Banfi, G. and Del, F. M. (2006). Serum creatinine values in elite athletes competing in 8 different sports: comparison with sedentary people. *Clinical Chemistry*, 52(2):330 -1.
8. Bisong, S., Brown, R. and Osim, E. (2011). Comparative effects of *Rauwolfia vomitoria* and chlorpromazine on social behaviour and pain. *North American Journal of Medical Sciences*, 3(1):48-54.
9. Branten, A. J., Vervoort, G. and Wetzels, J. F. (2005). Serum creatinine is a poor marker of GFR in nephrotic syndrome. *Nephrology Dialysis Transplantation*, 20(4):707-11.
10. Brien, C. B. (2009). The hospitalised patient with abnormal liver function tests. *Clin Liver Dis*, 13(2):179-92.
11. Cheng, F., Melissa, B., Fang, X., Shijun, L. Meghan, D., Hongxiu, L., Weizhu, Y., Kenneth, A, Xinxin, D. and Jun, G. (2008). Mechanism of chloroform-induced renal toxicity: Non-involvement of hepatic cytochrome P450-dependent metabolism. *Toxicology and Applied Pharmacology*, 227 (1): Pages 48-55.
12. Cholongitas, E., Shusang, V., Marelli, L., Nair, D., Thomas, M., Patch, D., Burns, A., Sweny, P. and Burroughs, A. K. (2007). Review article: renal function assessment

- in cirrhosis - difficulties and alternativemeasurements. *Alimentary Pharmacology & Therapeutics*,26(7):969-78.
13. Corbett, J. V. (2008). Laboratory tests and diagnostic procedures with nursing diagnoses; pp. 90-107.
 14. Dharnidharka, V. R., Kwon, C. and Stevens, G. (2002). Serum cystatin C is superior to serum creatinine as a marker of kidney function: a meta-analysis. *American Journal of Kidney Diseases*,40(2):221-6.
 15. Hawley, G. G. (2015). The Condensed Chemical Dictionary, 10th ed. Van Nostrand Reinhold, New York, NY, p. 237.
 16. Hughes, J. and Jefferson, A. (2008). Clinical chemistry made easy. 1st ed. New York, NY: Churchill Livingstone, p. 125-149.
 17. International Programme on Chemical Safety (IPCS) (2017).Chloroform. Environmental Health Criteria 163, WHO: Geneva.
 18. IRIS, Integrated Risk Information System(2001).Toxicological Review of Chloroform. Washington. Environmental Protection Agency.
 19. Isaac, L. J., Abah, G., Akpan, B., and Ekaette, I. U. (2013).Haematological properties of different breeds and sexes of rabbits (p.24-27).Proceedings of the 18th Annual Conference of Animal Science Association of Nigeria.
 20. Johnston, D. E. (2012).Special Considerations in Interpreting Liver Function Tests.*AmFam Physician*, 59(8):2223-30.
 21. Krier, M. and Ahmed, A. (2009).The asymptomatic outpatient with abnormal liver function tests.*Clin Liver Dis*.13(2):167-77.
 22. Krishnamoorthi, R. and Bai, V. R. (2015).Phytochemical analysis and antioxidant property of Premnalatifolia.*Open Science Journal of Pharmacy and Pharmacology*. 3(2):6-9.
 23. Kutalek, R. and Prince, A. (2007).African medical plants. In: Yaniv Z, U. Bachrach (eds). Handbook of medicinal plants.CBS publishers, New Delhi.
 24. Lipinski, B. and Pretorius, E. (2012). Novel pathway of iron induced blood coagulation: implications for diabetes mellitus and its complications. *Polish Archives of Medical*, 122(3): 115-122.
 25. Merck, M. (2012).Haematologic reference ranges. Mareck Veterinary Manual. Retrieved from <http://www.merckmanuals.com/>
 26. Mesembe, O. E., Ibanga, I. and Osim, E. E. (2004).The effects of fresh and thermoxidized palm oil diets on some haematological indices in the rat. *Nigerian Journal of Physiological Sciences*,19:86-91.
 27. Miller, W. G., Myers, G. L., Ashwood, E. R., Killeen, A. A., Wang, E., Thienpont, L. M. and Siekmann, L. (2005). Creatinine measurement: state of the art in accuracy and interlaboratory harmonization. *Archives of Pathology & Laboratory Medicine*.129(3):297-304.
 28. NseAbasi, N. E., Mary, E. W., Uduak, A. and Edem, E. A. (2014).Haematological Parameters and Factors Affecting Their Values.*Agricultural Science*, 2(1):37-47
 29. Ogunbajo, S. O., Alemode, I. C., Adama, J. Y. and Abdullahi, J. (2009).Haematological parameters of Savannah brown does fed varying dietary levels of flamboyant tree seed meal (p. 88-91). Proceedings of 34th Annual Conference of Nigerian Society for Animal roduction.
 30. Togun, V. A., Oseni, B. S. A., Ogundipe, J. A., Arewa, T. R., Hamed, A. A., Ajonijebu, D. C. and Mustapha, F. (2007). Effects of chronic lead administration on the haematological parameters of rabbits - a preliminary study (p. 341). Proceedings of the 41st Conferences of the Agricultural Society of Nigeria.

Ugwu *et al*

31. U.S. Environmental Protection Agency (U.S. EPA) (2011). Toxicological Review of Chloroform.
32. Ugwu, O. P. C., Edwin, N. and Ogbanshi, M. E. (2015). The Effect

www.iaajournals.org
of Ethanol Leaf Extract of
Jatropha curcas on Chloroform
Induced Hepatotoxicity in Albino
Rats. *Global Journal of
Biotechnology & Biochemistry* 10 (1):
11-15.